

Container label

MaxForce[®] Fly Spot Bait

GROUP	4	INSECTICIDE
-------	---	-------------

FOR INDOOR USE ONLY
For control of house flies

Only to be used by individuals holding an appropriate pesticide applicator certificate or license recognized by the provincial/territorial pesticide regulatory agency where the application occurs

COMMERCIAL

WETTABLE GRANULES

GUARANTEE: Imidacloprid.....10%
Muscalure.....0.1%

REGISTRATION NUMBER 32493 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 50 g to 0.5 Kg

BAYER CROPSCIENCE INC.
Suite 200, 160 Quarry Park Blvd SE.
Calgary, Alberta
T2C 3G3

Product information:
1-888-283-6847
www.bayeres.ca

In case of spills, poisoning or fire, telephone emergency response number (toll-free) 1-800-334-7577 (24 hours a day)

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

DO NOT APPLY IN THE PRESENCE OF CHILDREN OR IN AREAS ACCESSIBLE TO CHILDREN

Harmful if swallowed. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear a long-sleeved shirt, long pants, socks, shoes, and chemical-resistant gloves during application and clean-up activities.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

TOXICOLOGICAL INFORMATION: No specific antidote is available. Treat the patient symptomatically.

ENVIRONMENTAL HAZARDS: Toxic to aquatic organisms. Toxic to birds, do not spray birds. Toxic to bees, do not apply this product or allow it to drift if bees are present. Apply this product only in accordance with this label.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed. Store in original containers and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage.

DISPOSAL

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.

2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

MAXFORCE® is a registered trademark of the Bayer Group.

150819

Instruction booklet

MaxForce[®] Fly Spot Bait

GROUP	4	INSECTICIDE
-------	---	-------------

FOR INDOOR USE ONLY

For control of house flies

Only to be used by individuals holding an appropriate pesticide applicator certificate or license recognized by the provincial/territorial pesticide regulatory agency where the application occurs

COMMERCIAL

WETTABLE GRANULES

GUARANTEE: Imidacloprid.....10%
Muscalure.....0.1%

REGISTRATION NUMBER 32493 PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 50 g to 0.5 Kg

BAYER CROPSCIENCE INC.
Suite 200, 160 Quarry Park Blvd SE.
Calgary, Alberta
T2C 3G3

Product information:
1-888-283-6847
www.bayeres.ca

**In case of spills, poisoning or fire, telephone emergency response number
1-800-334-7577 (24 hours a day)**

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID: Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

TOXICOLOGICAL INFORMATION: No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. DO NOT APPLY IN THE PRESENCE OF CHILDREN OR IN AREAS ACCESSIBLE TO CHILDREN. Harmful if swallowed. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

PROTECTIVE CLOTHING AND EQUIPMENT: Wear a long-sleeved shirt, long pants, socks, shoes, and chemical resistant gloves during application and clean-up activities.

ENVIRONMENTAL HAZARDS: Toxic to aquatic organisms. Toxic to birds, do not spray birds. Toxic to bees, do not apply this product or allow it to drift if bees are present. Apply this product only in accordance with this label.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal.


STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed. Store in original containers and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage.

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.

4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

GENERAL INFORMATION

- A bait that is applied as a spray to be consumed by house flies.
- Contains a specially formulated mixture of attractants.
- Contains 2 powerful attractants designed to lure house flies to their death.
- Contains the attractant muscalure
- Contains Bitrex® bittering agent to help prevent ingestion by non-target animals, children, and pets.
- Easy to apply.
- Residual efficacy can last up to 4 weeks unless the bait is completely consumed.

<p>QUESTIONS ??</p> 	<p>For questions or comments, call toll-free 1-888-283-6847.</p>
--	--

DIRECTIONS FOR USE

Product Information

This product is bait formulated to control house flies in commercial and residential structures. The product is to be diluted and applied as a spray for treatment of structural surfaces and waste containers.

Use MaxForce Fly Spot Bait as part of an overall integrated pest management program for house flies which should include proper sanitation and environmental controls, such as exclusion and proper light management.

Applications of this product may leave a visible residue or discolor some surfaces. Before treating, it is advisable to test application in a small inconspicuous area.

TO CONTROL HOUSE FLIES	
<p>LOCATIONS OF USE</p>	<p>FOR INDOOR USE ONLY.</p> <p>For use in commercial and residential structures such as non-food/non-feed areas of commercial food handling establishments, restaurants, dining areas, storage facilities, schools, hospitals, nursing homes, offices and buildings at marinas. May also be applied inside dumpsters and other waste containers.</p> <p><u>Food/Feed Handling Establishments:</u></p> <p>Applications in food/feed handling establishments are permitted ONLY in non-food/non-feed areas. Non-food/non-feed areas include garbage rooms,</p>

	<p>lavatories, entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, mop closets and storage (after packaging, canning or bottling).</p> <p>Food/feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Included also are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food.</p> <p>Contamination of food or food contact surfaces must be avoided. Do not apply this product to any food, food packaging, or any food contact surfaces.</p> <p>Not for use in agricultural production facilities.</p>
WHERE TO APPLY	<p>Spray diluted MaxForce Fly Spot Bait on structural surfaces where house flies tend to rest such as on walls, around windows and the inside surfaces of dumpsters or trash collection bins.</p> <p>House flies prefer warm and sites with no air movement for resting. Avoid application in cool areas and in areas with air movement such as near fans.</p> <p>As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.</p> <p>DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.</p> <p>Treated baits spots must be placed in locations NOT accessible to children, pets, livestock or wildlife</p>
WHEN TO APPLY	Time applications to begin at the start of the season before fly populations have reached their peak.
DILUTION	<p>Do not apply undiluted.</p> <p>Add 125 g of MaxForce Fly Spot Bait/L of water. Agitate until completely mixed.</p>
USE RATE	<ul style="list-style-type: none"> • For low to moderate infestations of house flies, apply two litres of finished spray to treat 100 m² of fly-resting surfaces. • For moderate to high infestations of house flies, apply four litres of finished spray to treat 100 m² of fly-resting surfaces.
APPLICATION EQUIPMENT	<p><u>Spray Application:</u></p> <p>For spray applications, apply as a coarse fan or cone spray. Apply solution at a low pressure not to exceed 135 kPa (20 PSI). Flush spray equipment with water to prevent clogging.</p>
HOW TO APPLY	<p><u>Spray Application:</u></p> <p>There is no need to spray the complete surface because MaxForce Fly Spot Bait will attract house flies to the treated area. Best results occur when MaxForce Fly Spot Bait is applied to many locations throughout the structure or facility as a spot treatment. Apply product in spots ranging from 0.3 to 1.0 m² in locations where house flies tend to rest.</p>

RE-APPLICATION	Re-apply if house flies continue to be a problem, but not before 7 days.
RESTRICTIONS	<ul style="list-style-type: none"> • Do not apply bait on surfaces accessible to children or pets. • Do not allow bait to come in contact with food or water, livestock feed or their water. • Do not apply in food areas of food processing or manufacturing facilities. • Do not apply in classrooms or patient rooms. • Do not apply in agricultural production facilities. • Do not exceed 500 g of MaxForce Fly Bait per 100 m².

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that MaxForce Fly Spot Bait contains a Group 4 insecticide. Any insect population may contain individuals naturally resistant to MaxForce Fly Spot Bait and other Group 4 insecticides. The resistant individuals may dominate the insect population if this group of insecticides is used repeatedly in the same location. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of MaxForce Fly Spot Bait or other Group 4 insecticides with different groups that control the same pests.
- Insecticide use should be based on an Integrated Pest Management (IPM) program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information and to report suspected resistance, contact a Bayer CropScience representative at 1-888-283-6847 or at www.bayercropscience.ca.

150818