



Active Ingredients: Iprodione (23.1%) + Trifloxystrobin (1.44%) Equivalent to 256 g Iprodione and 56 g Trifloxystrobin per litre

FRAC Code: 2 + 11 (Dicarboximide + Qol)

Systemicity: localized penetrant **Formulation:** suspension concentrate

Signal Word: caution

Best Uses: Dollar spot, brown patch, snow moulds, Microdochium patch, leaf spots

Key Strengths: Control of DMI-resistant dollar spot, turfgrass summer safety, plant health promotion

Flexible disease control and stress management are important for turfgrass managers. Interface STRESSGARDTM provides flexible, effective disease control under all conditions and mitigates plant stress. The combination of iprodione, trifloxystrobin and Stressgard Formulation Technology provides a solution for diseases and plant stresses under hot, cool, wet or dry conditions. As a non-DMI fungicide, Interface can be applied throughout the year without possible harmful plant growth regulator effects, and helps to control DMI-resistant plant pathogens. Interface delivers effective control of key diseases, provides plant health benefits and improves turf colour and quality.

Attributes of Interface STRESSGARD

- Flexible, broad-spectrum disease control
- Multiple modes of action to manage fungicideresistant plant pathogens
- Stressgard Formulation Technology for plant health promotion effects, summer stress reduction, UV radiation management, and improved turf colour and density

Solutions for Tough Diseases

- Controls DMI-resistant dollar spot
- Can be used for summer dollar spot and brown patch control without negative DMI turf growth regulating effects
- Promotes plant health
- Controls spring and fall diseases like leaf spots and Microdochium patch
- Use Interface Stressgard alone or in combination with Triton STRESSGARD for pink and gray snow mould control

How to use Interface STRESSGARD

Apply Interface when conditions are favourable for disease as part of an integrated management program on greens, tees or fairways. Interface Stressgard can be tank-mixed with Signature STRESSGARD as part of a summer decline management program. For extended control of gray snow mould, tank-mix Interface with Triton STRESSGARD, Daconil Ultrex®, or Daconil 2787 at recommended labelled rates.

KEY DISEASES CONTROLLED	RATE (PER 100 M²)	APPLICATION INTERVAL*	
Dollar Spot	95 - 160 mL	14 - 21 days (greens & tees) 14 - 28 days (fairways & roughs)	
Brown Patch	95 - 160 mL	14 - 21 days (greens & tees) 14 - 28 days (fairways & roughs)	
Microdochium (Fusarium) Patch	128 - 160 mL	14 - 21 days	
Pink Snow Mould	128 - 160 mL	before winter snow cover	
Grey Snow Mould (suppression)	95 - 160 mL	before winter snow cover*	

USE RATE (G/100 M²)	COVERAGE PER CONTAINER (100 M²)	COVERAGE PER CONTAINER (ACRES)	G TRIFLOXYSTROBIN PER ACRE	G IPRODIONE PER ACRE	APPLICATIONS AT RATE (PER YEAR)
95 mL	105	2.59	81	988	5
128 mL	78	1.92	83	1333	3.5
140 mL	71	1.75	91	1462	2.5
160 mL	62.5	1.53	104	1673	2

Container size: 10 L

2010 Microdochium Patch Trials - Oregon State University - Brian McDonald









Heriatge G - 64 oz - 21 days

Interface - 160 mL - 21 days

Untreated

Photos courtesy of R. Golembiewski - Bayer.





Trial initiated on a Poa annua putting green in Corvallis, Oregon on Feb. 22. Three total fungicide applications made on 21 day intervals.



www.bayeres.ca





