

Active Ingredients: Iprodione (23.1%) + Trifloxystrobin (1.44%)
Equivalent to 256 g Iprodione and 56 g Trifloxystrobin per litre
FRAC Code: 2 + 11 (Dicarboximide + QoI)
Systemicity: Localized penetrant
Formulation: Suspension concentrate
Signal Word: Caution

Best Uses: dollar spot, Microdochium patch, brown patch, anthracnose, snow moulds

Key Strengths: controls DMI-resistant dollar spot, turfgrass summer safety, plant health promotion

Flexible disease control and stress management are important for turfgrass managers. Interface STRESSGARD™ provides flexible, effective disease control under all conditions and mitigates plant stress. The combination of iprodione, trifloxystrobin and Stressgard Formulation Technology provides a solution for diseases and plant stresses under hot, cool, wet or dry conditions. As a non-DMI fungicide, Interface can be applied throughout the year without harmful plant growth regulator effects, and helps control DMI-resistant plant pathogens. Interface delivers effective control of key diseases, provides plant health benefits and improves turf colour and quality.

Attributes of Interface STRESSGARD



- Flexible, broad-spectrum disease control
- Multiple modes of action to manage fungicide-resistant plant pathogens
- Stressgard Formulation Technology for plant health promotion effects, summer stress reduction, UV radiation management, and improved turf colour and density

Solutions for Tough Diseases

- Controls DMI-resistant dollar spot
- Can be used for summer dollar spot, anthracnose and brown patch control without negative DMI turf growth regulating effects
- Promotes plant health
- Controls spring and fall diseases like leaf spots and Microdochium patch
- Use Interface alone or in combination for pink and gray snow mould control in areas receiving < 90 days snow cover

How to use Interface STRESSGARD

Apply Interface as a preventive or early curative treatment when conditions are favourable for dollar spot and other diseases as part of an integrated management program on greens, tees or fairways. Interface can be tank-mixed with Signature STRESSGARD as part of a spring clean-up and summer decline management program. For extended control of gray pink snow mould, tank-mix Interface with Daconil Ultrex®, or Daconil 2787 at recommended labelled rates.




Preventive dollar spot on putting greens at Michigan State University. Interface-treated plots at 95 mL/100 M² every 14 days and 160 mL/100 M² every 21 days. Photos taken August 13. Photos: Dr. Rob Golembiewski, Bayer.

USE RATE (G/100 M ²)	COVERAGE PER CONTAINER (100 M ²)	COVERAGE PER CONTAINER (ACRES)	G TRIFLOXYSTROBIN PER ACRE	G IPRODIONE PER ACRE	APPLICATIONS AT RATE (PER YEAR)
95 mL	105	2.59	61	988	5
128 mL	78	1.92	83	1333	3.5
140 mL	71	1.75	91	1462	2.5
160 mL	62.5	1.53	104	1673	2

Container size: 10 L

KEY DISEASES CONTROLLED	RATE (PER 100 M ²)	APPLICATION INTERVAL*
Dollar Spot	95 - 160 mL	14 - 21 days (greens & tees) 14 - 28 days (fairways & roughs)
Anthracnose	95 - 160 mL	14 days (basal rot) 14 - 21 days (foliar)
Brown Patch	95 - 160 mL	14 - 21 days (greens & tees) 14 - 28 days (fairways & roughs)
Microdochium (Fusarium) Patch	128 - 160 mL	14 - 21 days
Pink Snow Mould	128 - 160 mL	before winter snow cover
Grey Snow Mould (suppression)	95 - 160 mL	before winter snow cover*

*See interface STRESSGARD label for a complete reference on control options and application instructions. Always read and carefully follow label instructions. All applications should be applied in sufficient water volume (6-8 L water/100 M²).



Preventive dollar spot trials on fairway turf at St. Thomas Golf and Country Club. All applications were made at indicated rates every 21 days starting on June 22, 2015. Data shown were taken during peak disease pressure on Aug. 26, 2015.

