



FLOREL

Maximize your plants appearance and production.


Fact Sheet

A foliar spray of FLOREL™ Plant Growth Regulator will increase lateral branching of many potted greenhouse grown ornamental species. Maximize your plants appearance and production. For use on stock plants, on rooting transplants and in final production.

Foliar applications of FLOREL™ Plant Growth Regulator will increase lateral branching of many potted greenhouse grown ornamental species thereby reducing stem elongation. This can eliminate manual pinching of some species.

Features and Benefits of Applying FLOREL

- ☑ Convenient Packaging for small or large applications (4 x 1 L Case).
- ☑ An ethylene-producing chemical that increases branching.
- ☑ Faster branching with little or no pinching.



HOW IT WORKS

When applied to plants, FLOREL Plant Growth Regulator readily enters the plant and breaks down to ethylene, a naturally occurring plant hormone. Ethylene production within the plant is stimulated by stress. For this reason, it is important that plants being treated are not under stress from drought, high temperature, disease or other environmental stresses.



FLOREL

Application Rate:

- ☑ New Guinea Impatiens - mix 104 mL of FLOREL Plant Growth Regulator in 100 L of water (250 ppm) and apply to thoroughly wet the foliage, but not to runoff.
- ☑ Begonia, chrysanthemum (perennial species), fuchsia, geranium, sweet potato vine (ipomea), lantana, poinsettia, verbena, and vinca vines (Vinca major) - mix 208 mL of FLOREL Plant Growth Regulator in 100 L of water (500 ppm) and apply to thoroughly wet the foliage, but not to runoff.
- ☑ Use low rate of FLOREL 104 mL / 100 L under low light conditions and on compact varieties as a strong growth regulating effect can occur under these conditions.
- ☑ Use of higher rate of FLOREL Plant Growth Regulator is recommended once experience has been gained with the product on specific crops.
- ☑ Apply FLOREL Plant Growth Regulator spray solution within 24 hours after mixing.

Application Timing:

- ☑ Stock Plants - Apply FLOREL Plant Growth Regulator to stock plants on a 14 day interval with a maximum of 4 applications. Apply in the evening, or early morning. Ensure a high soil temperature (21- 23°C) and a low carrier pH (5.5) is maintained. Make applications at normal pinching times instead of hand pinching. To optimize the vigour of cuttings, do not make applications for two weeks prior to harvesting cuttings from stock plants.
- ☑ Finished Plants - Apply FLOREL Plant Growth Regulator 14 to 21 days after transplanting of rooted cuttings. Re-apply every 10 to 14 days with a maximum of 4 applications. Apply in the evening, or early morning. Ensure a high soil temperature (21- 23°C) and a low carrier pH (5.5) is maintained.
- ☑ To ensure flowering and full foliage on finished plants, do not make applications for 6 to 8 weeks prior to bloom or planned sale.
- ☑ Not for use on cut flowers.
- ☑ Do not apply in more than 1 L per 10m² of bench space.
- ☑ DO NOT enter into treated areas during the restricted-entry interval (REI) of 12 hours.



Multiple Florel applications showing maximum branching.

Environmental Impact

All pesticides are regulated under the Pest Control Products Act to ensure that they do not pose an unacceptable risk to human health and the environment. For information regarding the pesticide regulatory process, visit Health Canada's (Pest Management Regulatory Agency) website www.hc-sc.gc.ca/pmra-arla/

